

University of Nebraska - Lincoln

DigitalCommons@University of Nebraska - Lincoln

---

Bulletin of the University of Nebraska State  
Museum

Museum, University of Nebraska State

---

11-1970

## Machairodont Cats from the Early Pleistocene Broadwater and Lisco Local Faunas

C. Bertrand Schultz

Larry D. Martin

Follow this and additional works at: <https://digitalcommons.unl.edu/museumbulletin>



Part of the [Entomology Commons](#), [Geology Commons](#), [Geomorphology Commons](#), [Other Ecology and Evolutionary Biology Commons](#), [Paleobiology Commons](#), [Paleontology Commons](#), and the [Sedimentology Commons](#)

---

This Article is brought to you for free and open access by the Museum, University of Nebraska State at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Bulletin of the University of Nebraska State Museum by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

BULLETIN OF  
*The University of Nebraska State Museum*

VOLUME 9, NUMBER 2  
NOVEMBER, 1970

C. Bertrand Schultz  
Larry D. Martin

# Machairodont Cats from the Early Pleistocene Broadwater and Lisco Local Faunas





Frontispiece.—View (1939) of a portion of the main Broadwater fossil quarry (University of Nebraska State Museum Collecting Locality Mo-5) located on the Dan J. Boman ranch,  $5\frac{1}{2}$  miles East and  $\frac{3}{4}$  mile North of Broadwater, Morrill County, Nebraska. The quarry was excavated from 1936 to 1942, including two winters, and as many as 38 men worked at one time at this site. It represents the most extensive single quarry excavation in the State of Nebraska. Additional test excavations also were made in 1965 and 1967. The holotype of *Ischyrosmilus crusafonti*, new species, was found in this quarry. The U.N.S.M. Coll. Loc. Mo-6 is located approximately  $\frac{1}{4}$  mile East of the Coll. Loc. Mo-5.

C. Bertrand Schultz

Larry D. Martin

**Machairodont Cats from  
the Early Pleistocene  
Broadwater and Lisco  
Local Faunas**

BULLETIN OF *The University of Nebraska State Museum*

VOLUME 9, NUMBER 2  
NOVEMBER, 1970

## ABSTRACT

### Machairodont Cats from the Early Pleistocene Broadwater and Lisco Local Faunas

C. Bertrand Schultz

Larry D. Martin

A new species of *Ischyrosmilus* (*I. crusafonti*) is reported from the Early Pleistocene Broadwater Local Fauna of Morrill County, Nebraska. *Ischyrosmilus* sp. is reported from the Early Pleistocene Lisco Local Fauna. *Megantereon* is reported from the Broadwater Local Fauna and this is the first record of the genus from North America.

# The Machairodont Cats from the Early Pleistocene Broadwater and Lisco Local Faunas.<sup>3</sup>

## INTRODUCTION

The Early Pleistocene Broadwater Formation from western Nebraska near Broadwater and Lisco in Morrill and Garden counties has yielded felid remains which should aid in a better understanding of machairodont cats.

The genus *Smilodon* has been reported from the Early Pleistocene deposits of Nebraska by Barbour and Schultz (1939, p. 20), Schultz and Stout (1941, p. 23, Table 3; 1948, p. 563, Table 2), and Schultz (1950, p. 68, Fig. 15) but these identifications were based on incomplete specimens, including the premaxilla with canines present (U.N.S.M. 1105), obtained from deposits of the Lisco Member of the Broadwater Formation. McGrew (1944, p. 55) lists *Smilodon* sp. from the Early Pleistocene, Sand Draw Local Fauna. This record was based on a femur, F. M. (Field Museum of Natural History) No. P26164,

lacking the distal end, but the present writers have not seen the specimen. However, McGrew's description, "The femur from Sand Draw differs from that of *Smilodon californicus* in having a straighter, longer, and more slender shaft and in having a deeper notch between the great trochanter and head," would seem to fit *Ischyrosmilus* better than *Smilodon*. It is now evident that *Smilodon* remains have not been recovered thus far from the Early Pleistocene deposits of Nebraska, and the saber-toothed felid examples from these sediments should be referred to the two genera *Ischyrosmilus* and *Megantereon*.

*Ischyrosmilus crusafonti*,<sup>4</sup> new species

*Holotype*.—Partial mandible with I<sub>1</sub>-/C, P<sub>3</sub> (alv.)-M<sub>1</sub>, U.N.S.M.<sup>5</sup> 25493 (Fig. 1, B).

*Referred Specimen from Type Locality*.—Partial left C/, U.N.S.M. 25503 (Fig. 1, C).

*Type Locality*.—U.N.S.M. Coll. Loc. Mo-5

<sup>1</sup> Director of the University of Nebraska State Museum and Regents Professor of Geology, Department of Geology, Lincoln.

<sup>2</sup> Research Assistant, Department of Geology and University of Nebraska State Museum (1969-1970); field party leader, Division of Vertebrate Paleontology of the Museum (1965-1969).

<sup>3</sup> The research for this paper has been supported by grants from the University of Nebraska Research Council, including a Research Assistantship (September, 1969, to June, 1970) for the junior author. The Board of Regents, University of Nebraska, also granted the senior author a leave of absence from his regular duties for six months (1969-1970) for paleontological research in Europe and U.S.A.

<sup>4</sup> Named in honor of Professor Miguel Crusafont Pairo, in recognition of his important work on European fossil felids, and for his scholarly work in connection with the Instituto Provincial de Paleontologia de Sabadell in north-eastern Spain.

<sup>5</sup> Abbreviations used in description: alv., alveolus or alveoli; br., broken; rt., root or roots; U.N.S.M., University of Nebraska State Museum; K.U.M.V.P., University of Kansas Museum of Natural History; U.C.M.P., University of California Museum of Paleontology; U.S.N.M., U.S. National Museum; W.T., West Texas State College, Panhandle Plains Historical Museum.

(NE.¼, Sec. 20, T. 19N., R.47W.) on the Dan Boman ranch, 5¼ mi. E. and ¾ mi. N. of Broadwater, Morrill County, Nebraska. A skeleton of *Stegomastodon* (mounted in the U.N.S.M. for exhibit) was found at Coll. Loc. Mo-5. The holotypes of *Procastoroides sweeti* Barbour and Schultz and *Satherium priscinaria middleswarti* Barbour and Schultz are from U.N.S.M. Coll. Loc. Mo-6, which is of the same geologic age and located only ¼ mi. E. of Coll. Loc. Mo-5 (Barbour and Schultz, 1937). (See Frontispiece.)

**Stratigraphic Occurrence.**—Early Pleistocene, Lisco Member, Broadwater Formation.

**Diagnosis.**—Smallest known species of genus;  $P_3$  single rooted, but proportionally large.

**Discussion.**—The mandible (U.N.S.M. 25493) is considerably smaller and more lightly built than the holotype (ramus) of *Ischyrosmilus ischyryus* Merriam, but it resembles the latter in most other respects. The dependent flange of *I. crusafonti* is developed to about the same degree as in *I. ischyryus*, and there are two large mental foramina along the posterior-ventral border of the flange and under  $P_3$ . These foramina are also similarly situated in the other species of the genus. The margin of the flange underneath the canine is vertical and the anterior portion of the ramus makes a flat rectangular surface. The ramus is proportionally shallower below the  $P_3$  than it is in *I. ischyryus*. The incisors of *I. crusafonti* are caniniform, each consisting of a single cusp which is recurved posteriorly. The incisors and the canine are serrated as in *Machairodus caticopus* Cope (Burt, 1931, Pl. 45). The  $I_1$  is small and compressed while  $I_2$  and  $I_3$  become progressively larger. The canine is relatively large for a machairodont cat. The  $P_3$  was single rooted as shown by the alveolus. The  $P_3$  appears to have been proportionally larger than that in *I. ischyryus*. The  $P_4$  of *I. crusafonti* is slightly recurved posteriorly and has two roots. It consists of a paraconid, a larger protoconid, and a metaconid about equal in size to the paraconid. An accessory cusp (talonid) also may have been present posterior to the metaconid, but the tooth is too worn to be certain of this. The  $M_1$  also is worn but shows a blade-like paraconid and protoconid with a deep carnassial notch between them.

The partial left C/ is coarsely serrated and, although the distal one-third is missing, the canine appears to have been a relatively short, broad tooth.

#### *Ischyrosmilus* sp.

**Material.**—Premaxilla with  $I^1$  (alv.)—C/, U.N.S.M. 1105 (Fig. 1, A).

**Locality.**—U.N.S.M. Coll. Loc. Gd-12, near center of E. ½, SE. ¼, Sec. 13, T. 18N., R. 46W. 3 mi. E., 2 mi. N. of Lisco, Garden County, Nebraska. This is the type locality of *Gigantocamelus fricki* Barbour and Schultz.

**Stratigraphic Occurrence.**—Early Pleistocene, Lisco Member, Broadwater Formation.

**Discussion.**—This felid was larger than *Ischyrosmilus crusafonti* from the Broadwater Local Fauna, and is near the size of *I. johnstoni* Mawby. The canines are proportionally shorter than in *Smilodon* although they are almost as broad. They are also more coarsely serrated. The incisors are arranged in a broad curve anterior to the canines, and are large and widely spaced. They show distinct wear facets in the regions where they interlocked with the lower incisors, as in *Smilodon*. The premaxilla is very heavy and massive.

#### *Megantereon hesperus* (Gazin)

**Referred Specimen.**—Partial mandible with  $I_2$ – $I_3$  alv., /C (br.),  $P_3$ – $P_4$  alv.,  $M_1$  U.N.S.M. 25494 (Fig. 2).

**Locality.**—U.N.S.M. Coll. Loc. Mo-5 (same locality as holotype of *Ischyrosmilus crusafonti*), Morrill County, Nebraska.

**Stratigraphic Occurrence.**—Early Pleistocene, Lisco Member, Broadwater Formation.

**Discussion.**—The holotype of *Megantereon hesperus* (Gazin) from the Early Pleistocene Hagerman Local Fauna of Idaho is a fragment of the posterior part of the ramus with  $M_1$ . The Nebraska mandible is more complete and demonstrates a pronounced dependent flange. There is a large mental foramen underneath the anterior margin of the alveolus for  $P_3$ , and a shallow but distinct groove slightly ventral to the midline of the lingual side of the ramus. The masseteric fossa extends forward to under the anterior root of  $M_1$  as it does in the holotype of




Fig. 1—A, U.N.S.M. 1105, *Ischyrosmilus* sp, premaxilla anterior and lateral views. B, U.N.S.M. 25493, holotype, *I. crusafonti* new species, left ramus, occlusal and lateral views. C, U.N.S.M. 25503, referred specimen, left upper canine. All X 3/5.



*M. hesperus* (Gazin). The mandible when viewed from the anterior end forms a broad rectangle with the flanges extending about 9 mm. below the ventral border of the symphysis on either side. There are several large foramina on both sides of the symphysis. The incisors are reduced and crowded. The left ramus shows alveolae for two small compressed incisors with  $I_1$  missing. The right ramus is probably partially pathologic and shows only the alveolus for  $I_3$ . The lower canine is proportionally large.  $P_3$  was double rooted as was  $P_4$ . Gazin (1933, p. 254) reported the following concerning the description of the holotype of *M. hesperus*: "The carnassial is relatively slender with the greatest transverse width well forward across the paraconid portion. The shearing blades . . . form a sharp angle when viewed from the side. Viewed from above the cutting edges form an angle which appears to be more acute than in *Machairodus catocopsis* . . . also less obtuse than in either *Smilodon gracilis* or *Smilodon californicus*." The Nebraska specimen agrees in all important respects.




Fig. 2—*Meganteron hesperus* (Gazin), referred, U.N.S.M. 25495, partial mandible (left ramus, lateral and lingual views; right ramus, lateral view), from the Lisco Member of the Broadwater Formation, Morrill County, Nebraska. X 3/5.

TABLE 1  
*Ischyrosmilus* MERRIAM  
COMPARATIVE MEASUREMENTS OF MANDIBULAR RAMI

	<i>Ischyrosmilus</i> <i>crusafonti</i> U.N.S.M. 25493 Holotype	<i>I. ischyros</i> U.C.M.P. 8140 Holotype	<i>I. johnstoni</i> <sup>1</sup> W.T. 1239 Holotype	<i>I. idahoensis</i> U.C.M.P. 22343 Holotype
Depth of flange.....	(53.6) <sup>2</sup>	59.	—	—
Depth of ramus anterior to $P_4$ .....	32.3	39.5	43.2	48.
Length of diastema.....	26.9	33.	40.	46.
Length $P_3$ - $M_1$ .....	(59.5)	(62.4)	69.8	(73.2)
Length $P_4$ - $M_1$ .....	43.8	(49.8)	48.9	58.5
Lower Canine anteroposterior.....	13.7	15.	—	18.
Lower Canine transverse.....	10.	12.5	—	13.5
$P_3$ anteroposterior .....	(10.)	7.5	13.0	(10.5)
$P_3$ transverse .....	(6.)	6.5	8.0	(6.)
$P_4$ anteroposterior .....	18.7	20.5	22.0	(24.)
$P_4$ transverse .....	9.6	12.5	10.3	(10.5)
$M_1$ anteroposterior .....	25.8	29.	28.5	(33.)
$M_1$ transverse .....	12.5	16.5	13.0	(13.)

<sup>1</sup> From Mawby, 1965, Table 1.

<sup>2</sup> ( ) = Approximate or measurement of alveolus.

TABLE 2  
*Megantereon hesperus* (GAZIN)  
 COMPARATIVE MEASUREMENTS OF MANDIBULAR RAMI

	U.N.S.M. 2549 Referred	U.S.N.M. 12614 Holotype
Depth of flange .....	(44.)	—
Depth of ramus posterior to M <sub>1</sub> .....	27.7	24.
Length of diastema .....	22.5	—
Length P <sub>3</sub> -M <sub>1</sub> .....	(49.)	—
Length P <sub>4</sub> -M <sub>1</sub> .....	(37.8)	—
Lower Canine anteroposterior .....	11.6	—
Lower Canine transverse .....	8.2	—
P <sub>3</sub> anteroposterior .....	10.2	—
P <sub>3</sub> transverse .....	(5.8)	—
P <sub>4</sub> anteroposterior .....	(18.2)	—
P <sub>4</sub> transverse .....	(7.2)	—
M <sub>1</sub> anteroposterior .....	19.7	19.8
M <sub>1</sub> transverse .....	9.3	8.5

## SUMMARY AND CONCLUSIONS

It seems likely that *Ischyrosmilus* is descended from *Machairodus* (*Heterofelis*) of the Pliocene of North America. The writers are aware of no valid records of *Machairodus* from the Early Pleistocene (Blancan) of North America. "*Machairodus*" *hesperus* Gazin is considered to be a species of *Megantereon* in this paper. The left P<sub>4</sub> (K.U.M.V.P. 3917) from the Rexroad Local Fauna assigned to *Machairodus* sp. by Hibbard (1937, p. 243, Pl. 2, Fig. 13) also appears to belong to the genus *Megantereon*. The anteroposterior diameter for the latter tooth is 18.5 mm. and this compares closely with the alveolar measurement of 18.2 mm. for the P<sub>4</sub> (U.N.S.M. 25494) of *M. hesperus*. The Kansas specimen (K.U.M.V.P. 3917) is also very similar to the P<sub>4</sub> of *Megantereon megantereon* illustrated by Viret (1954, Pl. 13, Fig. 3).

Mawby (1965, p. 584) compared *Ischyrosmilus* with *Homotherium crenatidens* Fabrini from the Early Pleistocene (Villafranchian) of Europe and suggests that they might be congeneric. This is possible since the skulls of the two forms are quite similar. However, in *H. crenatidens* the dependent flange is not so well developed on the ramus and its anterior border is inclined rather than being vertical as in *Ischyrosmilus*. Considering the differences in the anterior portion of the mandible, it would appear that the two forms developed independently from a common generic stock (?*Machairodus*). In this case the genus *Ischyrosmilus* Merriam should be retained, and probably also *Dinobastis* Cope. *Ischyrosmilus crusafonti*, new species, is the

fourth species of the genus to be recognized. All of these species are Blancan in age, and until some idea of size variation can be obtained it remains questionable if they are all valid. However, *I. crusafonti* is decidedly distinct from the other three species.

The similarity between the Blancan faunas of North America and the Villafranchian faunas of Eurasia is very striking, and with further study many more forms will undoubtedly be found to occur in both faunas. This will cause some difficulty as more and more names will be found to be preoccupied by names from the other continent. For instance, *Megantereon hesperus* (Gazin) is doubtfully separate from *Megantereon megantereon* (Croizet and Jobert), although they probably should not be synonymized until better material of *M. hesperus* can be compared. *Megantereon* apparently immigrated from Eurasia to North America at the beginning of the Pleistocene. The extinction of the Barbourufelini at the end of the upper Pliocene (Kimballian) would have left open the niche for dirk-toothed felids in North America. *Megantereon* may then have given rise to *Smilodon*. The earliest known examples of *Smilodon* appear to be those from the late Yarmouthian or early Illinoian deposits in Sheridan County, Nebraska.

## ACKNOWLEDGMENTS

The present writers are indebted to the following for assistance and critical comment: Professors T. M. Stout and Harvey L. Gunderson, and Messrs. Lloyd G. Tanner and King A. Richey. Thanks are due to Prof. Donald E.

Savage for the loan of felid types from the collections of the University of California Museum of Paleontology. The illustrations were prepared by Jerry Tanner and the manuscript was typed by Norma Wagner and Mary Cutler. The Dan J. Boman and Mary A. Boman families kindly granted permission to the University of Nebraska State Museum to excavate for fossils on their properties near Broadwater, and Mr. Frank Ruby assisted the Museum in obtaining permission to excavate at the localities near Lisco. The U.N.S.M. fossils used in the present report were collected by field parties, which included the following members: David Abbott, E. L. Blue, Paul Burkholder, Frank Crabill, Gordon Graham, Jack Graham, Joe Johnson, Dean Kerl, Fred Schall, Marian and Bertrand Schultz, Thompson M. Stout, and Lloyd G. Tanner. The field parties were aided by local collectors from Bridgeport, namely Messrs. W. F. Chaloupka, T. C. Middleswart, John Ochoa, and S. R. Sweet, and also by more than twenty workers furnished by the Works Progress Administration during 1938-1942. The writers are grateful to Dr. Clayton Ray of the U.S. National Museum for the use of the holotype of *Meganteron hesperus* (Gazin).

## REFERENCES

- Barbour, Erwin H., and C. Bertrand Schultz. 1937. An Early Pleistocene fauna from Nebraska. *Amer. Mus. Novitates* (942): 1-10, Figs. 1-4.
- . 1939. A new giant camel. *Bull. Univ. Nebraska State Mus.* 2 (2): 17-27, Figs. 5-12.
- Churcher, C. S. 1965. The affinities of *Dinobastes serus* Cope 1893. Abstracts 7th Congress International Assoc. Quaternary Research (1965), Boulder: 65.
- Chardin, Pierre Teilhard de. 1945. Les felides de Chine. Les formes fossiles: 3-36, 20 figures *in* Chardin and P. Leroy, Publ. Inst. Geobiol. 11: 1-58, 21 figures.
- Gazin, C. L. 1933. New felids from the Upper Pliocene of Idaho. *Jour. Mammal.*, 14: 251-256, 3 figs.
- Hay, O. P. 1930. Second bibliography and catalogue of the fossil vertebrata of North America. Carnegie Inst. Washington, Publ. (390), Vol. 2: i-xiv, 1-1074.
- Hubbard, Claude W. 1937. An upper Pliocene fauna from Meade County, Kansas. *Trans. Kansas Acad. Sci.* 40: 239-263, Figs. 1-2, Pls. 1-5.
- Kurten, Bjorn. 1965. The Pleistocene Felidae of Florida. *Bull. Florida State Mus., Biol. Sci.* 9 (6): 215-273, Figs. 1-17, 31 tables.
- Matthew, W. D. 1918. Contributions to the Snake Creek fauna, with notes upon the Pleistocene of western Nebraska, American Museum expedition of 1916. *Bull. Amer. Mus. Nat. Hist.* 38 (7): 183-229, Figs. 1-20, Pls. 4-10.
- Mawby, John E. 1965. Machairodonts from the Late Cenozoic of the Panhandle of Texas. *Jour. Mammal.* 46 (4): 573-587, Figs. 1-5.
- McGrew, Paul O. 1944. An Early Pleistocene (Blancan) fauna from Nebraska. *Field Mus. Nat. Hist. Geol. Ser.* 9 (2), Publ. 546: 33-66, Figs. 14-22.
- Meade, Grayson E. 1961. The saber-toothed cat, *Dinobastis serus*. *Bull. Texas Mem. Mus.* (2): 23-60.
- Merriam, John C. 1905. A new sabre-tooth from California. *Univ. California Publ. Bull. Dept. Geol. Sci.* 4 (9): 171-175, Fig. 1.
- . 1918. New mammalia from the Idaho Formation. *Ibid.*, 10 (26): 523-530, Figs. 1-5.
- . 1919. Tertiary mammalian faunas of the Mohave Desert. *Ibid.*, 11 (5): 437a-437e, 438-585, Figs. 1-253.
- Merriam, John C., and Chester Stock. 1932. The Felidae of Rancho La Brea. Carnegie Inst. Wash. Publ. (422): 1-231, Figs. 1-152, 42 plates, 114 tables.
- Schultz, C. Bertrand. 1950. Stratigraphic distribution of Pleistocene mammals in Nebraska. *In* Condra, G. E., E. C. Reed, and E. D. Gordon (Revised by G. E. Condra and E. C. Reed), Correlation of the Pleistocene deposits of Nebraska. *Bull. Nebraska Geological Survey* (15A): 1-73, Figs. 1-12, 2 plates.
- Schultz, C. Bertrand, Marian R. Schultz, and Larry D. Martin. 1970. A new tribe of saber-toothed cats (Barbourofelini) from the Pliocene of North America. *Bull. Univ. Nebraska State Mus.* 9 (1): in press.
- Schultz, C. Bertrand, and Thompson M. Stout. 1941. Guide for field conference on the Tertiary and Pleistocene of Nebraska. Special Publ. Univ. of Nebraska State Mus., (1): 1-55, Figs. 1-47, 2 charts. 1 map.
- . 1945. Pleistocene loess deposits of Nebraska. *Amer. Jour. Sci.* 243: 231-244, Figs. 1-5, Pls. 1-2.
- . 1948. Pleistocene mammals and terraces in the Great Plains. *Bull. Geol. Soc. Amer.* 59 (6): 553-558, Figs. 1-4, Pl. 1, 3 tables.
- Stout, Thompson M. 1965. Broadwater Formation. *In* Schultz and Smith, Editors, *Guidebook for (INQUA) Field Conference D, Central Great Plains*, VIIth Congress, International Assoc. Quaternary Research, Nebraska Academy of Sciences Publ., Lincoln: 67-68, 1 fig.
- Viret, M. J. 1954. Le loess a bancs durcis de Saint-Vallier (Drome) et sa faune de mammiferes Villafranchiens. *Nouv. Arch. Mus. d'Hist. Nat., Lyon*, 4: 1-200, 43 figs., 33 pls.

#### **THE BOARD OF REGENTS**

Richard E. Adkins, Osmond  
J. G. Elliott, Scottsbluff  
B. N. Greenberg, M.D., York  
Richard L. Herman, Omaha  
Robert L. Raun, Minden  
Edward Schwartzkopf, Lincoln  
G. Robert Ross, Lincoln, Corporation Secretary

#### **THE CHANCELLOR**

D. B. Varner

#### **THE PRESIDENT, LINCOLN CAMPUSES AND OUTSTATE ACTIVITIES**

Joseph Soshnik